

A Generalization of Power’s Archimedean Circles

Hiroshi Okumura and Masayuki Watanabe

Abstract. We generalize the Archimedean circles in an arbelos given by Frank Power.

Let three semicircles α , β and γ form an arbelos with inner semicircles α and β with diameters PA and PB respectively. Let a and b be the radii of the circles α and β . Circles with radii $t = \frac{ab}{a+b}$ are called Archimedean circles. Frank Power [2] has shown that for “highest” points Q_1 and Q_2 of α and β respectively, the circles touching γ and the line OQ_1 (respectively OQ_2) at Q_1 (respectively Q_2) are Archimedean (see Figure 1). We generalize these Archimedean circles.




Figure 1

We denote the center of γ by O . Let Q be the intersection of the circle γ and the perpendicular of AB through P , and let δ be a circle touching γ at the point Q from the inside of γ . The radius of δ is expressed by $k(a+b)$ for a real number k satisfying $0 \leq k < 1$. The tangents of δ perpendicular to AB intersect α and β at points Q_1 and Q_2 respectively, and intersect the line AB at points P_1 and P_2 respectively (see Figures 2 and 3).

Theorem. (1) *The radii of the circles touching the circle γ and the line OQ_1 (respectively OQ_2) at the point Q_1 (respectively Q_2) are $2(1-k)t$.*
 (2) *The circle touching the circles γ and α at points different from A and the line P_1Q_1 from the opposite side of B and the circle touching the circles γ and β at points different from B and the line P_2Q_2 from the opposite side of A are congruent with common radii $(1-k)t$.*




Figure 2

Proof. (1) Since $|PP_1| = 2ka$, $|OP_1| = (b - a) + 2ka$. While

$$|P_1Q_1|^2 = |PP_1||P_1A| = 2ka(2a - 2ka) = 4k(1 - k)a^2.$$

Hence $|OQ_1|^2 = ((b - a) + 2ka)^2 + 4k(1 - k)a^2 = (a - b)^2 + 4kab$. Let x be the radius of one of the circles touching γ and the line OQ_1 at Q_1 . From the right triangle formed by O , Q_1 and the center of this circle, we get

$$(a + b - x)^2 = x^2 + (a - b)^2 + 4kab$$

Solving the equation for x , we get $x = \frac{2(1-k)ab}{a+b} = 2(1 - k)t$. The other case can be proved similarly.




Figure 3

(2) The radius of the circle touching α externally and γ internally is proportional to the distance between the center of this circle and the radical axis of α and γ [1, p. 108]. Hence its radius is $(1 - k)$ times of the radii of the twin circles of Archimedes. \square

The Archimedean circles of Power are obtained when δ is the circle with a diameter OQ . The twin circles with the half the size of the Archimedean circles in [4] are also obtained in this case. The statement (2) is a generalization of the twin circles of Archimedes, which are obtained when δ is the point circle. In this case the points Q_1 , Q_2 and P coincide, and we get the circle with radius $2t$ touching the line AB at P and the circle γ by (1) [3].

References

- [1] J. L. Coolidge, *A treatise on the circle and the sphere*, Chelsea. New York, 1971 (reprint of 1916 edition).
- [2] Frank Power, Some more Archimedean circles in the arbelos, *Forum Geom.*, 5 (2005) 133–134.
- [3] H. Okumura and M. Watanabe, The twin circles of Archimedes in a skewed arbelos, *Forum Geom.*, 4 (2004) 229–251.
- [4] H. Okumura and M. Watanabe, Non-Archimedean twin circles in the arbelos,(in Bulgarian), *Math. Plus*, 13 (2005) no.1, 60–62.

Hiroshi Okumura: Department of Information Engineering, Maebashi Institute of Technology,
460-1 Kamisadori Maebashi Gunma 371-0816, Japan
E-mail address: okumura@maebashi-it.ac.jp

Masayuki Watanabe: Department of Information Engineering, Maebashi Institute of Technology,
460-1 Kamisadori Maebashi Gunma 371-0816, Japan
E-mail address: watanabe@maebashi-it.ac.jp