

## Circle Chains Inside a Circular Segment

Giovanni Lucca

**Abstract.** We consider a generic circles chain that can be drawn inside a circular segment and we show some geometric properties related to the chain itself. We also give recursive and non recursive formulas for calculating the centers coordinates and the radius of the circles.

### 1. Introduction

Consider a circle with diameter  $AB$ , center  $C$ , and a chord  $GH$  perpendicular to  $AB$  (see Figure 1). Point  $O$  is the intersection between the diameter and the chord. Inside the circular segment bounded by the chord  $GH$  and the arc  $GBH$ , it is possible to construct a doubly infinite chain of circles each tangent to the chord, and to its two immediate neighbors.




Figure 1. Circle chain inside a circular segment

Let  $2(a + b)$  be the diameter of the circle and  $2b$  the length of the segment  $OB$ . We set up a cartesian coordinate system with origin at  $O$ . Beginning with a circle with center  $(X_0, Y_0)$  and radius  $r_0$  tangent to the chord  $GH$  and the arc  $GBH$ , we construct a doubly infinite chain of tangent circles, with centers  $(X_i, Y_i)$  and radius  $r_i$  for integer values of  $i$ , positive and negative.

## 2. Some geometric properties of the chain

We first demonstrate some basic properties of the doubly infinite chain of circles.

**Proposition 1.** *The centers of the circles lie on the parabola with axis along  $AB$ , focus at  $C$ , and vertex the midpoint of  $OB$ .*




Figure 2. Centers of circles in chain on a parabola

*Proof.* Consider a circle of the chain with center  $O'(x, y)$ , radius  $r$ , tangent to the arc  $GBH$  at  $Q$ . Since the segment  $CQ$  contains  $O'$  (see Figure 2), we have, by taking into account that  $C$  has coordinates  $(b - a, 0)$  and

$$\begin{aligned} CQ &= a + b, \\ CO' &= \sqrt{(x - b + a)^2 + y^2}, \\ O'Q &= r = x, \\ CO' &= CQ - O'Q. \end{aligned}$$

From these, we have

$$\sqrt{(x - b + a)^2 + y^2} = a + b - x,$$

which simplifies into

$$y^2 = -4a(x - b). \quad (1)$$

This clearly represents the parabola symmetric with respect to the  $x$ -axis, vertex  $(b, 0)$ , the midpoint of  $OB$ , and focus  $(b - a, 0)$ , which is the center  $C$  of the given circle.  $\square$

**Proposition 2.** *The points of tangency between consecutive circles of the chain lie on the circle with center  $A$  and radius  $AG$ .*




Figure 3. Points of tangency on a circular arc

*Proof.* Consider two neighboring circles with centers  $(X_i, Y_i)$ ,  $(X_{i+1}, Y_{i+1})$ , radii  $r_i, r_{i+1}$  respectively, tangent to each other at  $T_i$  (see Figure 3). By using Proposition 1 and noting that  $A$  has coordinates  $(-2a, 0)$ , we have

$$AO_i^2 = (X_i + 2a)^2 + Y_i^2 = \left(-\frac{Y_i^2}{4a} + b + 2a\right)^2 + Y_i^2,$$

$$r_i^2 = X_i^2 = \left(-\frac{Y_i^2}{4a} + b\right)^2.$$

Applying the Pythagorean theorem to the right triangle  $AO_iT_i$ , we have

$$AT_i^2 = AO_i^2 - r_i^2 = 4a(a + b) = AO \cdot AB = AG^2.$$

It follows that  $T_i$  lies on the circle with center  $A$  and radius  $AG$ . □

**Proposition 3.** *If a circle of the chain touches the chord  $GH$  at  $P$  and the arc  $GBH$  at  $Q$ , then the points  $A, P, Q$  are collinear.*

*Proof.* Suppose the circle has center  $O'$ . It touches  $GH$  at  $P$  and the arc  $GBH$  at  $Q$  (see Figure 4). Note that triangles  $CAQ$  and  $O'PQ$  are isosceles triangles with  $\angle ACQ = \angle PO'Q$ . It follows that  $\angle CQA = \angle O'QP$ , and the points  $A, P, Q$  are collinear. □




Figure 4. Line joining points of tangency

*Remark.* Proposition 3 gives an easy construction of the circle given any one of the points of tangency. The center of the circle is the intersection of the line  $CQ$  and the perpendicular to  $GH$  at  $P$ .

### 3. Coordinates of centers and radii

Figure 5 shows a right triangle  $O_i O_{i-1} K_i$  with the centers  $O_{i-1}$  and  $O_i$  of two neighboring circles of the chain. Since these circles have radii  $r_{i-1} = X_{i-1}$  and  $r_i = X_i$  respectively, we have

$$\begin{aligned} (X_{i-1} - X_i)^2 + (Y_i - Y_{i-1})^2 &= (r_i + r_{i-1})^2 = (X_i + X_{i-1})^2, \\ (Y_i - Y_{i-1})^2 &= 4X_i X_{i-1}. \end{aligned}$$

Making use of (1), we rewrite this as

$$(Y_i - Y_{i-1})^2 = 4 \left( b - \frac{Y_i^2}{4a} \right) \left( b - \frac{Y_{i-1}^2}{4a} \right),$$

or

$$\frac{4a(a+b) - Y_{i-1}^2}{4a^2} \cdot Y_i^2 - 2Y_{i-1}Y_i + \frac{(a+b)Y_{i-1}^2 - 4ab^2}{a} = 0. \quad (2)$$

If we index the circles in the chain in such a way that the ordinate  $Y_i$  increases with the index  $i$ , then from (2) we have

$$Y_i = \frac{2Y_{i-1} - \left( \frac{Y_{i-1}^2}{a} - 4b \right) \sqrt{1 + \frac{b}{a}}}{2 \left( 1 + \frac{b}{a} - \frac{Y_{i-1}^2}{4a^2} \right)}. \quad (3a)$$

This is a recursive formula that can be applied provided that the ordinate  $Y_0$  of the first circle is known. Note that  $Y_0$  must be chosen in the interval  $(-2\sqrt{ab}, 2\sqrt{ab})$ .



This is to distinguish from the extension of the chain by working the recursion (3a) backward:<sup>1</sup>

$$Y_{i-1} = \frac{2Y_i + \left(\frac{Y_i^2}{a} - 4b\right) \sqrt{1 + \frac{b}{a}}}{2 \left(1 + \frac{b}{a} - \frac{Y_i^2}{4a^2}\right)}. \tag{3b}$$

Thus, for negative integer values of  $i$ , with

$$Z_{-i} = \frac{Y_{-i}}{2a} + \sqrt{1 + \frac{b}{a}},$$

we have

$$Z_{-i} = -\frac{1}{-\alpha - \frac{1}{-\alpha - \frac{1}{\dots - \frac{1}{-\alpha + Z_{0-}}}}},$$

where

$$Z_{0-} = \frac{Y_0}{2a} + \sqrt{1 + \frac{b}{a}}.$$

It is possible to give nonrecursive formulas for calculating  $Y_i$  and  $Y_{-i}$ . For brevity, in the following, we shall consider only  $Y_i$  for positive integer indices because, as far as  $Y_{-i}$  is concerned, it is enough to change, in all the formulae involved,  $\alpha$  into  $-\alpha$ ,  $Z_i$  into  $Z_{-i}$ , and  $Z_{0+}$  into  $Z_{0-}$ . Starting from (5), and by considering its particular structure, one can write, for  $i = 1, 2, 3, \dots$ ,

$$Z_i = -\frac{Q_{i-1}(\alpha)}{Q_i(\alpha)}$$

where  $Q_i(\alpha)$  are polynomials with integer coefficients. Here are the first ten of them.

$Q_0(\alpha)$	1
$Q_1(\alpha)$	$\alpha + Z_{0+}$
$Q_2(\alpha)$	$(\alpha^2 - 1) + \alpha Z_{0+}$
$Q_3(\alpha)$	$(\alpha^3 - 2\alpha) + (\alpha^2 - 1)Z_{0+}$
$Q_4(\alpha)$	$(\alpha^4 - 3\alpha^2 + 1) + (\alpha^3 - 2\alpha)Z_{0+}$
$Q_5(\alpha)$	$(\alpha^5 - 4\alpha^3 + 3\alpha) + (\alpha^4 - 3\alpha^2 + 1)Z_{0+}$
$Q_6(\alpha)$	$(\alpha^6 - 5\alpha^4 + 6\alpha^2 - 1) + (\alpha^5 - 4\alpha^3 + 3\alpha)Z_{0+}$
$Q_7(\alpha)$	$(\alpha^7 - 6\alpha^5 + 10\alpha^3 - 4\alpha) + (\alpha^6 - 5\alpha^4 + 6\alpha^2 - 1)Z_{0+}$
$Q_8(\alpha)$	$(\alpha^8 - 7\alpha^6 + 15\alpha^4 - 10\alpha^2 + 1) + (\alpha^7 - 6\alpha^5 + 10\alpha^3 - 4\alpha)Z_{0+}$
$Q_9(\alpha)$	$(\alpha^9 - 8\alpha^7 + 21\alpha^5 - 20\alpha^3 + 5\alpha) + (\alpha^8 - 7\alpha^6 + 15\alpha^4 - 10\alpha^2 + 1)Z_{0+}$

According to a fundamental property of continued fractions [1], these polynomials satisfy the second order linear recurrence

$$Q_i(\alpha) = \alpha Q_{i-1}(\alpha) - Q_{i-2}(\alpha). \tag{7}$$

<sup>1</sup>Equation (3b) can be obtained by solving equation (2) for  $Y_{i-1}$ .

We can further write

$$Q_i(\alpha) = P_i(\alpha) + P_{i-1}(\alpha)Z_{0+}, \quad (8)$$

for a sequence of simpler polynomials  $P_i(\alpha)$ , each either odd or even. In fact, from (7) and (8), we have

$$P_{i+2}(\alpha) = \alpha P_{i+1}(\alpha) - P_i(\alpha).$$

Explicitly,

$$P_i(\alpha) = \begin{cases} 1, & i = 0, \\ \sum_{k=0}^{\frac{i}{2}} (-1)^{\frac{i}{2}+k} \binom{\frac{i}{2}+k}{2k} \alpha^{2k}, & i = 2, 4, 6, \dots, \\ \sum_{k=1}^{\frac{i+1}{2}} (-1)^{\frac{i+1}{2}+k} \binom{\frac{i-1}{2}+k}{2k-1} \alpha^{2k-1}, & i = 1, 3, 5, \dots \end{cases}$$

From (6), we have

$$Y_i = a \left( \alpha - 2 \frac{Q_{i-1}(\alpha)}{Q_i(\alpha)} \right),$$

for  $i = 1, 2, \dots$

## References

- [1] H. Davenport, *Higher Arithmetic*, 6-th edition, Cambridge University Press, 1992.

Giovanni Lucca: Via Corvi 20, 29100 Piacenza, Italy  
*E-mail address:* vanni\_lucca@inwind.it