

## Some Circles Associated with the Feuerbach Points

Nguyen Thanh Dung

**Abstract.** Consider a triangle with its nine-point circle tangent to the incircle and excircles at the Feuerbach point. We show that the four circles each through the circumcenter, nine-point, and Feuerbach point contain the nine-point center of the intouch triangle or the corresponding extouch triangle. Furthermore, the lines joining these Feuerbach points to the corresponding nine-point centers are concurrent on the nine-point circle of the given triangle.

### 1. Four coaxial circles through the Feuerbach points

The starting point of this note is the famous Feuerbach theorem, that for a given triangle, the nine-point circle is tangent internally to the incircle and externally to each of the excircles. Given triangle  $ABC$ , with incenter  $I$ , excenters  $I_a, I_b, I_c$ , and nine-point  $N$ , the points of tangency of the nine-point circle with these circles are the Feuerbach points  $F_e, F_a, F_b, F_c$  on the lines  $NI, NI_a, NI_b, NI_c$  with ratios of division

$$NF_e : F_e I = \frac{R}{2} : -r,$$

$$NF_a : F_a I_a = \frac{R}{2} : r_a, \quad NF_b : F_b I_b = \frac{R}{2} : r_b, \quad NF_c : F_c I_c = \frac{R}{2} : r_c,$$

where  $R, r, r_a, r_b, r_c$  are the circumradius, inradius, and exradii.

**Proposition 1.** *Let  $O$  be the circumcenter of triangle  $ABC$ .*

(a)  $OI^2 = R(R - 2r)$ .

(b)  $OI_a^2 = R(R + 2r_a)$ .

(c)  $OI_a^2 = 4R(r_a - r)$ .

(d) *The excentral triangle  $I_a I_b I_c$  has circumcenter  $I'$  at the reflection of  $I$  in  $O$ , and circumradius  $2R$ .*

For (a-c), see [1, Theorems 152-154]. For (d), see [4, §4.6.1].

Consider the circle through  $O, N$ , and  $F_e$ . Since  $I$  is an interior point of the segment  $NF_e$ , it is an interior point of the circle. Our first result relates the endpoint of the chord through  $O$  and  $I$  with the intouch triangle, whose vertices are the points of tangency of the incircle with the sidelines.

**Theorem 2.** *The line  $OI$  intersects the circle  $ONF_e$  again at the nine-point center of the intouch triangle.*




Figure 1

*Proof.* Let  $N_i$  be the second intersection of the line  $OI$  with the circle  $ONF_e$ . By the intersecting chords theorem,  $OI \cdot IN_i = NI \cdot IF_e$ . Therefore,

$$\frac{IN_i}{OI} = \frac{NI \cdot IF_e}{OI^2} = \frac{\left(\frac{R}{2} - r\right) r}{R(R - 2r)} = \frac{r}{2R}.$$

Note that the intouch triangle is homothetic to the excentral triangle. Since the excentral triangle has circumcenter  $I'$  and nine-point center  $O$ , its Euler line is the line  $OI$ . Since the intouch triangle has circumcenter  $I$ , its Euler line is also the line  $OI$ . From

$$\frac{IN_i}{I'O} = \frac{IN_i}{OI} = \frac{r}{2R},$$

the homothetic ratio of the intouch and excentral triangles, we conclude that  $N_i$  is nine-point center of the intouch triangle.  $\square$




Figure 2.

Analogous results hold if we replace the Feuerbach point  $F_e$  by the other Feuerbach points, say,  $F_a$ . if the circle through  $O, N, F_a$  intersects the line  $OI_a$  at  $N_a$ , then  $\frac{I_a N_a}{I_a O} = \frac{r_a}{2R}$ . Now, triangle  $I I_b I_c$  is homothetic to the  $A$ -extouch triangle formed by the points of tangency of the  $A$ -excircle with the sidelines of

triangle  $ABC$ , with homothetic ratio  $-\frac{r}{2R}$ , since the circumradius of  $II_bI_c$  is also  $2R$ . In fact, the circumcenter of  $II_bI_c$  is the reflection of  $I_a$  in  $O$ . It follows that  $\frac{I_a N_a}{I_a O} = -\frac{I_a N_a}{O I_a} = -\frac{r}{2R}$ , and  $N_a$  is the nine-point center of the  $A$ -extouch triangle (see Figure 2).

**Theorem 3.** *The points  $O, N, F_a$  and  $N_a$  are concyclic; so are  $O, N, F_b, N_b$ , and  $O, N, F_c, N_c$ .*

**2. Concurrency of four lines on the nine-point circle**

In preparation for the proof of our next main result (Theorem 7 below), we establish an interesting relation between the centers  $O, N, I, I_a$  given in Proposition 5. The reformulation as Proposition 6 in terms of directed angles ([2, §§16-19]) makes the proof of Theorem 7 independent of the relative position of  $O$  and  $N$  with respect to the bisector of angle  $A$ .

**Lemma 4.** *Let  $N$  be the nine-point center of triangle  $ABC$ .*

- (a) *If  $A > 60^\circ$ , then  $N$  and  $O$  lie on the same side of the bisector of angle  $A$ .*
- (b) *If  $A = 60^\circ$ , then  $N$  lies on the bisector of angle  $A$ .*
- (c) *If  $A < 60^\circ$ , then  $N$  and  $O$  lie on opposite sides of the bisector of angle  $A$ .*




Figure 3A




Figure 3B

*Proof.* First consider the case when  $A$  is an obtuse angle (see Figure 3A). Construct the perpendicular from  $O$  to  $BC$ , to intersect the circumcircle at  $M$  on the opposite side of  $A$ . The line  $AM$  is clearly the bisector of angle  $A$ . If  $X$  is the orthogonal projection of  $A$  on  $BC$ , then the orthocenter  $H$  and  $X$  are on opposite sides of  $A$ . It follows that  $O$  and  $H$ , and their midpoint  $N$ , all are on the same side of the bisector  $AM$ .

The same conclusion holds if  $A = 90^\circ$ , since the orthocenter  $H$  coincides with  $A$ .

Now we assume  $A$  an acute angle (see Figure 3B). It is known that  $AH = 2R \cos A$ , and that the bisector of angle  $A$  also bisects the angle  $OAH$ . It divides

$OH$  in the ratio  $R : 2R \cos A = 1 : 2 \cos A$ . Therefore,  $O$  and  $N$  are on the same side of the bisector if and only if  $2 \cos A < 1$ , i.e.,  $A > 60^\circ$ .

If  $A = 60^\circ$ , then  $AH = AO$ , and  $N$  lies on the bisector. This completes the proof of the theorem.  $\square$

**Proposition 5.** (a) *The angle  $\angle IOI_a$  is acute, right, or obtuse according as  $A$  is less than, equal to, or greater than  $60^\circ$ .*

$$(b) \angle INI_a = \begin{cases} 2\angle IOI_a, & \text{if } A \leq 60^\circ, \\ 360^\circ - 2\angle IOI_a, & \text{if } A > 60^\circ. \end{cases}$$

*Proof.* (a) Applying the law of cosines to triangle  $IOI_a$ , and using the expressions for the lengths given in Proposition 2, we have

$$\begin{aligned} \cos \angle IOI_a &= \frac{OI^2 + OI_a^2 - II_a^2}{2 \cdot OI \cdot OI_a} \\ &= \frac{R(R - 2r) + R(R + 2r_a) - 4R(r_a - r)}{2 \cdot OI \cdot OI_a} \\ &= \frac{R(R - r_a + r)}{OI \cdot OI_a}. \end{aligned} \quad (1)$$

The angle  $\angle IOI_a$  is acute, right, or obtuse according as  $r_a - r$  is less than, equal to or greater than  $R$ . Since  $\frac{r_a - r}{R} = 4 \sin^2 \frac{A}{2}$ , these are the cases according as  $A$  is less than, equal to, greater than  $60^\circ$ .

(b) From (1), we have

$$\begin{aligned} \cos 2 \cdot \angle IOI_a &= 2 \cos^2 \angle IOI_a - 1 \\ &= \frac{(OI^2 + OI_a^2 - II_a^2)^2 - 2 \cdot OI^2 \cdot OI_a^2}{2 \cdot OI^2 \cdot OI_a^2} \\ &= \frac{(2R(R - r_a + r))^2 - 2R(R - 2r) \cdot R(R + 2r_a)}{2R(R - 2r) \cdot R(R + 2r_a)} \\ &= \frac{R^2 - 6R(r_a - r) + 2(r_a^2 + r^2)}{(R - 2r)(R + 2r_a)}. \end{aligned} \quad (2)$$

On the other hand,

$$\begin{aligned} \cos \angle INI_a &= \frac{NI^2 + NI_a^2 - II_a^2}{2NI \cdot NI_a} \\ &= \frac{\left(\frac{R}{2} - r\right)^2 + \left(\frac{R}{2} + r_a\right)^2 - 4R(r_a - r)}{2\left(\frac{R}{2} - r\right)\left(\frac{R}{2} + r_a\right)} \\ &= \frac{R^2 - 6R(r_a - r) + 2(r_a^2 + r^2)}{(R - 2r)(R + 2r_a)}. \end{aligned} \quad (3)$$

Comparison of (2) and (3) shows that  $\cos \angle INI_a = \cos 2 \cdot \angle IOI_a$ . Therefore,  $\angle INI_a = 2\angle IOI_a$  or  $360^\circ - 2\angle IOI_a$ . Taking (a) into account, we have (b).  $\square$

*Remark.* When  $A = 60^\circ$ ,  $\angle IOI_a = 90^\circ$ , and  $\angle INI_a = 180^\circ$  (see Lemma 4(b)).

In terms of directed angles, we reformulate Proposition 5 below.

**Proposition 6.**  $(NI, NI_a) = -2(OI, OI_a)$ .

**Theorem 7.** *The four lines  $F_eN_i$ ,  $F_aN_a$ ,  $F_bN_b$ ,  $F_cN_c$  are concurrent at a point on the nine-point circle.*

*Proof.* It is enough to prove this for the lines  $F_eN_i$  and  $F_aN_a$  (see Figure 4). Let  $P$  be the intersection of the line  $F_eN_i$  with the nine-point circle. We show that it also lies on the line  $F_aN_a$ . For this, it is enough to verify  $(F_aN, F_aP) = (F_aN, F_aN_a)$ .




Figure 4.

$$\begin{aligned}
 (F_aN, F_aP) &= (PF_a, PN) && \text{triangle } NF_aP \text{ isosceles} \\
 &= (PF_a, PF_e) + (PF_e, PN) \\
 &= \frac{1}{2}(NF_a, NF_e) + (PF_e, PN) && N = \text{center of circle } PF_eF_a \\
 &= \frac{1}{2}(NI_a, NI) + (PF_e, PN) \\
 &= (OI, OI_a) + (PF_e, PN) && \text{Proposition 6} \\
 &= (OI, OI_a) + (NF_e, N_iF_e) && \text{triangle } NPF_e \text{ isosceles} \\
 &= (OI, OI_a) + (NO, N_iO) && O, N, F_e, N_i \text{ concyclic} \\
 &= (OI, ON_a) + (ON, OI) \\
 &= (ON, ON_a) \\
 &= (F_aN, F_aN_a) && O, N, F_a, N_a \text{ concyclic.}
 \end{aligned}$$

The same reasoning shows that  $P$  also lies on the lines  $F_bN_b$  and  $F_cN_c$ . □

We summarize the main results in this note in Figure 5 below, and conclude with an identification of the triangle centers  $N_i$  and  $P$ . According to the ENCYCLOPEDIA OF TRIANGLE CENTERS [3], the intouch triangle has orthocenter  $X(65)$ . Its nine-point center  $N_i$ , being the midpoint of  $IX(65)$ , is  $X(942)$ . This point lies on the line through  $X(11) = F_e$  and  $X(113)$ , which is on the nine-point circle. Therefore  $P$  is  $X(113)$ , which is also the midpoint of  $HX(110)$ .




Figure 5.

## References

- [1] N. A. Court, *College Geometry*, Dover reprint, 2007.
- [2] R. A. Johnson, *Advanced Euclidean Geometry*, Dover reprint, 2007.
- [3] C. Kimberling, *Encyclopedia of Triangle Centers*, available at <http://faculty.evansville.edu/ck6/encyclopedia/ETC.html>.
- [4] P. Yiu, *Introduction to the Geometry of the Triangle*, Florida Atlantic University Lecture Notes, 2001; with corrections, 2013, available at <http://math.fau.edu/Yiu/Geometry.html>

Nguyen Thanh Dung: Chu Van An high school for Gifted students, 55 To Son street, Chi Lang ward, Lang son province, Viet Nam

*E-mail address:* nguyenthanhdungcva@gmail.com