Forum Geometricorum
Volume 18 (2018) 103–114.

FORUM GEOM
ISSN 1534-1178

Geometric Inequalities in Pedal Quadrilaterals

Şahlar Meherrem, Gizem Günel Açıksöz, Serenay Şen, Zeynep Sezer, and Güneş Başkes

Abstract. The aim of this paper is to investigate the general properties of the pedal quadrilateral of a point P with respect to a convex and closed quadrilateral $ABCD$. In particular, we present an analogue of Erdős-Mordell Inequality, stating that for any triangle ABC and a point P inside ABC, the sum of the distances from P to the sides is less than or equal to half of the sum of the distances from P to the vertices of ABC, for an inscribed quadrilateral.

1. Introduction

Let ABC be a triangle and P be a point in plane. If the feet of the perpendiculars drawn from P to sides of the triangle $[BC]$, $[CA]$, $[AB]$ are respectively X, Y and Z, then triangle XYZ is called the pedal triangle of P with respect to ABC and P is called the pedal point ([3, 7]). If the point P lies on the circumcircle of the triangle ABC, then the points X, Y and Z are collinear. The line passing through the points X, Y and Z is known as the *Simson line* of P ([3]).



Figure 1

A similar notion has been generalized for polygons with n sides ([3]). Let $A_1A_2A_3\cdots A_n$ be a polygon and P be a point inside $A_1A_2A_3\cdots A_n$. If the feet of the perpendiculars from P to the line segments $[A_1A_2]$, $[A_2A_3]$, $\ldots$, $[A_nA_1]$ are respectively H_1, H_2, $\ldots$, H_n, then the polygon $H_1H_2\cdots H_n$ is called the *pedal polygon* of P with respect to $A_1A_2A_3\cdots A_n$ and P is called the pedal point ([3]). Although there are lots of investigations on pedal triangles, so little investigations have been studied on pedal quadrilaterals in the literature ([3, 5, 6, 7, 9, 11]).

Publication Date: April 20, 2018. Communicating Editor: Paul Yiu.

In this paper, we study on the general properties of the pedal quadrilateral of a point P with respect to a convex and closed quadrilateral $ABCD$. We investigate some interesting geometric inequalities on pedal quadrilaterals. In particular, we state an analogue of Erdős-Mordell Inequality ([1, 2, 4, 8, 10]) for quadrilaterals.

2. Geometric inequalities in pedal quadrilaterals

Proposition 1. *Let $ABCD$ be an inscribed quadrilateral and P be a point inside $ABCD$. If the feet of perpendiculars from P to the line segments $[AB]$, $[BC]$, $[CD]$ and $[AD]$ are respectively K, M, N, L and the radius of the circumcircle of $ABCD$ is R then,*

$$|KM| = \tfrac{|AC|\cdot|PB|}{2R}, \qquad |KL| = \tfrac{|BD|\cdot|PA|}{2R},$$

$$|MN| = \tfrac{|BD|\cdot|PC|}{2R}, \qquad |LN| = \tfrac{|AC|\cdot|PD|}{2R}.$$



Figure 2

Proof. As seen in Figure 2 and since $\angle BKP = \angle PLD = \angle PNC = \angle PMB = 90°$, the quadrilaterals $BKPM$, $PLDN$, $PNCM$ and $PLAK$ are inscribed quadrilaterals. By the law of sines, we have

$$|PB| = \frac{KM}{\sin KPM} \quad \text{and} \quad 2R = \frac{|AC|}{\sin(\pi - \angle KPM)} = \frac{|AC|}{\sin KPM}.$$

Hence, we get

$$|KM| = \frac{|AC|\cdot|PB|}{2R}.$$

It can be easily seen that

$$|KL| = \frac{|BD|\cdot|PA|}{2R}, \qquad |MN| = \frac{|BD|\cdot|PC|}{2R}, \qquad |LN| = \frac{|AC|\cdot|PD|}{2R}.$$

□

Proposition 2. *Let $ABCD$ be a quadrilateral and P be a point inside $ABCD$.If the feet of perpendiculars from P to the line segments $[AB]$, $[BC]$, $[CD]$ and $[AD]$ are respectively A_1, B_1, C_1, D_1, then*

$$r_1 + r_2 + r_3 + r_4 \leq \frac{M \cdot C}{2\sqrt{2}},$$

where r_1, r_2, r_3 and r_4 are respectively the radii of the incircles of the triangles AA_1D_1, A_1BB_1, CB_1C_1, DC_1D_1, and

$$M = \max\left\{\frac{1}{\sqrt{1-\cos A}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos B}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos C}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos D}+\sqrt{2}}\right\},$$

and C is the perimeter of the quadrilateral $ABCD$. The equality holds if $ABCD$ is chosen as a square and P is chosen at the center of the square.



Figure 3

Proof. By the law of cosines, we have

$$|A_1B_1| = \sqrt{|A_1B|^2 + |BB_1|^2 - 2|A_1B||BB_1|\cos B}. \tag{1}$$

If $S_{A_1BB_1}$ is the area of the triangle A_1BB_1, then

$$S_{A_1BB_1} = \frac{|A_1B| \cdot |BB_1| \sin B}{2}. \tag{2}$$

Since $2S_{A_1BB_1} = (|A_1B| + |BB_1| + |A_1B_1|)r_2$, we get

$$r_2 = \frac{2S_{A_1BB_1}}{|A_1B| + |BB_1| + |A_1B_1|}. \tag{3}$$

By the equalities (2) and (3),

$$r_2 = \frac{|A_1B| \cdot |BB_1| \sin B}{|A_1B| + |BB_1| + |A_1B_1|}. \tag{4}$$

It follows from Arithmetic-Geometric Mean Inequality that

$$|A_1B|^2 + |BB_1|^2 \geq 2 \cdot |A_1B| \cdot |BB_1|. \tag{5}$$

If the inequalities (1) and (5) are considered together, then

$$\begin{aligned} |A_1B_1| &= \sqrt{|A_1B|^2+|BB_1|^2-2|A_1B|\cdot|BB_1|\cos B} \\ &\geq \sqrt{2|A_1B|\cdot|BB_1|-2|A_1B|\cdot|BB_1|\cos B} \\ &\geq \sqrt{2}\,\sqrt{|A_1B|\cdot|BB_1|}\sqrt{1-\cos B} \end{aligned} \tag{6}$$

It is also obvious from Arithmetic-Geometric Mean Inequality that

$$|A_1B|+|BB_1| \geq 2\sqrt{|A_1B|\cdot|BB_1|}. \tag{7}$$

Therefore, by the inequalities (6) and (7), we observe that

$$|A_1B|+|BB_1|+|A_1B_1| \geq 2\sqrt{|A_1B|\cdot|BB_1|}+\sqrt{2}\,\sqrt{|A_1B|\cdot|BB_1|}\sqrt{1-\cos B}. \tag{8}$$

Hence, it follows from the inequalities (4) and (8) that

$$\begin{aligned} r_2 &\leq |A_1B|\cdot|BB_1|\frac{\sin B}{\sqrt{2}\cdot\sqrt{|A_1B|\cdot|BB_1|}\sqrt{1-\cos B}+2\sqrt{|A_1B|\cdot|BB_1|}} \\ &\leq |A_1B|\cdot|BB_1|\cdot\frac{\sin B}{\sqrt{2}\sqrt{|A_1B|\cdot|BB_1|}\sqrt{1-\cos B}+\sqrt{2})} \\ &\leq \sqrt{|A_1B|\cdot|BB_1|}\cdot\frac{\sin B}{\sqrt{2}(\sqrt{1-\cos B}+\sqrt{2})} \\ &\leq \frac{|A_1B|+|BB_1|}{2\sqrt{2}(\sqrt{1-\cos B}+\sqrt{2})} \end{aligned} \tag{9}$$

Let $M = \max\left\{\frac{1}{\sqrt{1-\cos A}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos B}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos C}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos D}+\sqrt{2}}\right\}$. By the inequality (9),

$$r_2 \leq \frac{M\cdot(|A_1B|+|BB_1|)}{2\sqrt{2}}. \tag{10}$$

Similarly, we get

$$\left.\begin{aligned} r_1 &\leq \tfrac{M\cdot(|A_1A|+|AD_1|)}{2\sqrt{2}}, \\ r_3 &\leq \tfrac{M\cdot(|C_1C|+|CB_1|)}{2\sqrt{2}}, \\ r_4 &\leq \tfrac{M\cdot(|C_1D|+|DD_1|)}{2\sqrt{2}}. \end{aligned}\right\} \tag{11}$$

If the inequalities (10) and (11) are added side by side, it is easy to see that

$$r_1+r_2+r_3+r_4 \leq \frac{M\cdot C}{2\sqrt{2}}$$

where C is the perimeter of the quadrilateral $ABCD$.

Now suppose that $ABCD$ is a square with side length $2a$ and P is at the center of the $ABCD$ (Figure 4). If the area of the triangle D_1AA_1 is $S_{D_1AA_1}$ then,

$$S_{D_1AA_1} = \frac{2a+a\sqrt{2}}{2}r_1 = \frac{a^2}{2}.$$



Figure 4

Hence, we get $r_1 = \frac{a}{2+\sqrt{2}}$. Since $r_1 = r_2 = r_3 = r_4$, $C = 8a$ and $M = \frac{1}{\sqrt{2}+1}$, then the following equality holds:

$$r_1 + r_2 + r_3 + r_4 = 4 \cdot \frac{a}{2+\sqrt{2}} = \frac{8a \cdot \left(\frac{1}{\sqrt{2}+1}\right)}{2\sqrt{2}} = \frac{C \cdot M}{2\sqrt{2}}.$$

□

Proposition 3. *Let $A_1A_2A_3 \cdot A_n$ be a polygon and P be a point inside $A_1A_2A_3 \cdots A_n$. If the feet of perpendiculars from P to the line segments $[A_1A_2]$, $[A_2A_3]$, …, $[A_nA_1]$ are respectively H_1, H_2, …, H_n, then*

$$r_1 + r_2 + \cdots + r_n \leq \frac{M \cdot C}{2\sqrt{2}}$$

where r_1, r_2, …, r_n are respectively the radii of the incircles of the triangles $A_1H_1H_n$, $H_1A_2H_2$, …, $H_{n-1}A_nH_n$, and

$$M = \max\left\{\frac{1}{\sqrt{1-\cos A_1}+\sqrt{2}}, \frac{1}{\sqrt{1-\cos A_2}+\sqrt{2}}, \ldots, \frac{1}{\sqrt{1-\cos A_n}+\sqrt{2}}\right\}$$

and C is the perimeter of the polygon $A_1A_2 \cdots A_n$. The equality holds if $A_1A_2 \cdots A_n$ is chosen as a square and P is chosen as the centroid of the square.

This can be easily shown as in Proposition 2.

Proposition 4. *Let $ABCD$ be a circumscribed quadrilateral with area S. If the center and the radius of the incircle of $ABCD$ are respectively the point P and r, and the feet of perpendiculars from P to the line segments $[AB]$, $[BC]$, $[CD]$ and*

$[AD]$ are respectively K, L, M and N, then

$$\frac{S_{\text{pedal}}}{S} \leq 1 - \frac{1}{4r}\sqrt{a^2+b^2+c^2+d^2},$$

where $a = |BL|$, $b = |LC|$, $c = |MD|$, $d = |AN|$, and S_{pedal} is the area of the pedal quadrilateral $KLMN$. The equality holds if $ABCD$ is a square.



Figure 5

Proof. As seen in Figure 5, if S_1, S_2, S_3 and S_4 are respectively the areas of triangles KPL, LPM, MPN and NPK, then we get

$$\begin{aligned}
S_1 &= a \cdot r - \frac{a^2 \sin B}{2},\\
S_2 &= b \cdot r - \frac{b^2 \sin C}{2},\\
S_3 &= c \cdot r - \frac{c^2 \sin D}{2},\\
S_4 &= d \cdot r - \frac{d^2 \sin A}{2}.
\end{aligned}$$

Hence, it is obvious that

$$S_1+S_2+S_3+S_4 = r(a+b+c+d) - \frac{1}{2}(a^2\sin B + b^2 \sin C + c^2 \sin D + d^2 \sin A).$$

Since we also have $S = (a+b+c+d)r$, we get

$$\begin{aligned}
\frac{S_{\text{pedal}}}{S} &= \frac{S_1+S_2+S_3+S_4}{S}\\
&= 1 - \frac{a^2 \sin B + b^2 \sin C + c^2 \sin D + d^2 \ \sin A}{2r(a+b+c+d)}. \qquad (12)
\end{aligned}$$

By the Cauchy-Schwarz Inequality, we get

$$\begin{aligned} a+b+c+d &= a\cdot 1+b\cdot 1+c\cdot 1+d\cdot 1 \\ &\le \sqrt{a^2+b^2+c^2+d^2}\cdot\sqrt{1^2+1^2+1^2+1^2} \\ &= 2\sqrt{a^2+b^2+c^2+d^2}. \end{aligned}$$

Thus,

$$-\frac{1}{a+b+c+d} \le -\frac{1}{2\sqrt{a^2+b^2+c^2+d^2}}.$$

By the equality (12),

$$\begin{aligned} \frac{S_{\text{pedal}}}{S} &= 1-\frac{a^2\sin B+b^2\sin C+c^2\sin D+d^2\sin A}{2r(a+b+c+d)} \\ &\le 1-\frac{a^2+b^2+c^2+d^2}{4r\sqrt{a^2+b^2+c^2+d^2}} \\ &= 1-\frac{1}{4r}\sqrt{a^2+b^2+c^2+d^2}. \end{aligned}$$



Figure 6

Now suppose that $ABCD$ is a square and $|AB| = |BC| = |CD| = |AD| = 2a$. As seen in Figure 6, since $r = a$, $S = 4a^2$, $S_{\text{pedal}} = 2a^2$, we have

$$\frac{S_{\text{pedal}}}{S} = \frac{2a^2}{4a^2} = \frac{1}{2} = 1-\frac{1}{4a}\sqrt{a^2+a^2+a^2+a^2} = 1-\frac{1}{4r}\sqrt{a^2+b^2+c^2+d^2}.$$

□

Proposition 5. *Let P be a point inside a polygon $A_1A_2A_3\cdots A_n$ where $|A_1A_2| = a_1$, $|A_2A_3| = a_2, \ldots, |A_nA_1| = a_n$. If the lengths of the perpendiculars from P to the sides of the polygon $A_1A_2A_3\cdots A_n$ are respectively $h_1, h_2, \ldots, h_n$, then*

$$\frac{S}{C^2} \ge \frac{1}{2}\cdot\frac{1}{\frac{a_1}{h_1}+\cdots+\frac{a_n}{h_n}}$$

where S and C are respectively the area and the perimeter of $A_1A_2A_3\cdots A_n$. The equality holds if $a_1 = a_2 = \cdots = a_n = a$ and $h_1 = h_2 = \cdots = h_n = h$.



Figure 7

Proof. As seen in Figure 7, it is obvious that $2S = a_1h_1 + a_2h_2 + \cdots + a_nh_n$. By the Cauchy-Schwarz Inequality, we get

$$(a_1h_1+a_2h_2+\cdots+a_nh_n)\left(\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}\right) \geq (\sqrt{a_1^2}+\sqrt{a_2^2}+\cdots+\sqrt{a_n^2})^2.$$

Since

$$2S\left(\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}\right) \geq (\sqrt{a_1^2}+\sqrt{a_2^2}+\cdots+\sqrt{a_n^2})^2$$

we have

$$2S \geq \frac{(a_1+a_2+\cdots+a_n)^2}{\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}} = \frac{C^2}{\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}}.$$

Hence,

$$\frac{S}{C^2} \geq \frac{1}{2}\cdot\frac{1}{\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}}.$$

Now suppose that $a_1 = a_2 = \cdots = a_n = a$, and $h_1 = h_2 = \cdots = h_n = h$. Since,

$$S = n\cdot\frac{ah}{2}, \qquad C = na,$$

we get

$$\frac{S}{C^2} = \frac{nah}{2n^2a^2} = \frac{h}{2na} = \frac{1}{2}\cdot\frac{1}{\frac{a}{h}+\frac{a}{h}+\cdots+\frac{a}{h}} = \frac{1}{2}\cdot\frac{1}{\frac{a_1}{h_1}+\frac{a_2}{h_2}+\cdots+\frac{a_n}{h_n}}.$$

□

Proposition 6. *Let P be a point inside a polygon $A_1A_2A_3\cdots A_n$ where $|A_1A_2| = a_1$, $|A_2A_3| = a_2, \ldots, |A_nA_1| = a_n$. If the lengths of the perpendiculars from P to the sides of the polygon $A_1A_2A_3\cdots A_n$ are respectively $h_1, h_2, \ldots, h_n$, then*

$$\frac{1}{a_1h_1} + \frac{1}{a_2h_2} + \cdots + \frac{1}{a_nh_n} \geq \frac{n^2}{2S}$$

where S is the area of $A_1A_2A_3\cdots A_n$. The equality holds if $a_1 = a_2 = \cdots = a_n = a$ and $h_1 = h_2 = \cdots = h_n = h$.

Proof. By the Cauchy-Schwarz Inequality, we have

$$(a_1h_1+a_2h_2+\cdots+a_nh_n)\left(\frac{1}{a_1h_1} + \frac{1}{a_2h_2} + \cdots + \frac{1}{a_nh_n}\right) \geq (\underbrace{1+1+\cdots+1}_{n \text{ times}})^2.$$

Since $2S = a_1h_1 + a_2h_2 + \cdots + a_nh_n$, it is easy to see that

$$\frac{1}{a_1h_1} + \frac{1}{a_2h_2} + \cdots + \frac{1}{a_nh_n} \geq \frac{n^2}{2S}.$$

Now suppose that $a_1 = a_2 = \cdots = a_n = a$ and $h_1 = h_2 = \cdots = h_n = h$. Since $S = n \cdot \frac{ah}{2}$, we have

$$\frac{1}{a_1h_1} + \frac{1}{a_2h_2} + \cdots + \frac{1}{a_nh_n} = n \cdot \frac{1}{ah} = \frac{n^2}{nah} = \frac{n^2}{2S}.$$

□

Theorem 7. *Let $ABCD$ be an inscribed quadrilateral and P be a point inside $ABCD$. If the feet of perpendiculars from P to the line segments $[AB]$, $[BC]$, $[CD]$ and $[AD]$ are A_1, B_1, C_1, D_1 respectively, then*

$$|PA| + |PB| + |PC| + |PD| > 4\sqrt[4]{|PA_1| \cdot |PA_2| \cdot |PA_3| \cdot |PA_4|}.$$



Figure 8

Proof. Let $|PA_1| = h_1$, $|PA_2| = h_2$, $|PA_3| = h_3$ and $|PA_4| = h_4$. Then by the law of cosines, we have

$$\begin{aligned}
|A_1A_2| &= \sqrt{h_1^2 + h_2^2 - 2h_1h_2\cos(\pi - B)},\\
|A_2A_3| &= \sqrt{h_2^2 + h_3^2 - 2h_2h_3\cos(\pi - C)},\\
|A_3A_4| &= \sqrt{h_3^2 + h_4^2 - 2h_3h_4\cos(\pi - D)},\\
|A_4A_1| &= \sqrt{h_4^2 + h_1^2 - 2h_1h_4\cos(\pi - A)}.
\end{aligned}$$

Hence,

$$\begin{aligned}
|A_1A_2|^2 &= h_1^2 + h_2^2 - 2h_1h_2\cos(\pi - B)\\
&= h_1^2 + h_2^2 - 2h_1h_2\cos(A + C + D - \pi)\\
&= h_1^2 + h_2^2 + 2h_1h_2\cos(A + C + D)\\
&= h_1^2 + h_2^2 + 2h_1h_2\cos(A + C)\cos D - 2h_1h_2\sin(A + C)\sin D\\
&= (h_2\sin D - h_1\sin(A + C))^2 + (h_2\cos D + h_1\cos(A + C))^2.
\end{aligned}$$

Since $A + C = \pi$, we have $|A_1A_2|^2 = (h_2\sin D)^2 + (h_2\cos D - h_1)^2$.

Now suppose that $(h_2\cos D - h_1)^2 = 0$. Then, $\cos D = \frac{h_1}{h_2}$. Since $\angle A_1PA_2 = \angle D$, by the law of cosines,

$$\begin{aligned}
|A_1A_2|^2 &= h_1^2 + h_2^2 - 2h_1h_2\cos D\\
&= h_1^2 + h_2^2 - 2h_1h_2 \cdot \frac{h_1}{h_2}\\
&= h_2^2 - h_1^2.
\end{aligned}$$

Hence, we get $h_2^2 = |A_1A_2|^2 + h_1^2$, implying that $\angle PA_1A_2 = 90^\circ$, which is a contradiction. Therefore, $(h_2\cos D - h_1)^2 \neq 0$. Then, it is easy to verify that

$$|A_1A_2| > h_2\sin D \tag{13}$$

Similarly, we have

$$|A_2A_3| > h_3\sin A \tag{14}$$

$$|A_3A_4| > h_4\sin B \tag{15}$$

$$|A_1A_4| > h_1\sin C \tag{16}$$

Since PA_1BA_2, PA_2CA_3, PA_3DA_4 and PA_4AA_1 are inscribed quadrilaterals, if we apply the law of sines to the triangles AA_1A_4, BA_1A_2, CA_2A_3, DA_3A_4,

then we respectively get,

$$|PA| = \frac{|A_1A_4|}{\sin A} \tag{17}$$

$$|PB| = \frac{|A_1A_2|}{\sin B} \tag{18}$$

$$|PC| = \frac{|A_2A_3|}{\sin C} \tag{19}$$

$$|PD| = \frac{|A_3A_4|}{\sin D} \tag{20}$$

If the inequalities (13), (14), (15) and (16) are respectively considered together with the inequalities (17), (18), (19) and (20), then

$$\begin{aligned}
|PB| &= \frac{|A_1A_2|}{\sin B} > \frac{h_2 \sin D}{\sin B},\\
|PC| &= \frac{|A_2A_3|}{\sin C} > \frac{h_3 \sin A}{\sin C},\\
|PD| &= \frac{|A_3A_4|}{\sin D} > \frac{h_4 \sin B}{\sin D},\\
|PA| &= \frac{|A_1A_4|}{\sin A} > \frac{h_1 \sin C}{\sin A}.
\end{aligned}$$

Therefore, it is easy to see that

$$\begin{aligned}
|PA| + |PB| + |PC| + |PD| &> \frac{h_1 \sin C}{\sin A} + \frac{h_2 \sin D}{\sin B} + \frac{h_3 \sin A}{\sin C} + \frac{h_4 \sin B}{\sin D}\\
&> 4\sqrt[4]{\frac{h_1 \sin C}{\sin A} \cdot \frac{h_2 \sin D}{\sin B} \cdot \frac{h_3 \sin A}{\sin C} \cdot \frac{h_4 \sin B}{\sin D}}\\
&= 4\sqrt[4]{|PA_1| \cdot |PA_2| \cdot |PA_3| \cdot |PA_4|}
\end{aligned}$$

by Arithmetic-Geometric Mean Inequality. □

References

[1] C. Alsina and R. B. Nelsen, A visual proof of the Erdős-Mordell inequality, *Forum Geom.*, 7 (2007) 99–102.

[2] L. Bankoff, An elementary proof of the Erdős-Mordell theorem, *Amer. Math. Monthly*, 65 (1958), 521.

[3] H. S. M. Coxeter and S. L. Greitzer, *Geometry Revisited*, Math. Assoc. America, 1967.

[4] P. Erdős, Problem 3740, *Amer. Math. Monthly*, 42 (1935) 396.

[5] D. Ferrarello, M. F. Mammana, and M. Pennisi, Pedal poygons, *Forum Geom.*, 13 (2013) 153–164.

[6] R. Honsberger, *Episodes In Nineteenth and Twentieth Century Euclidean Geometry*, Math. Assoc. America, 1995.

[7] R. A. Johnson, *Advanced Euclidean Geometry*, Dover Publications, 2007.

[8] D. K. Kazarinoff, A simple proof of the Erdős-Mordell inequality for triangles, *Michigan Math. Journal*, 4 (1957) 97–98.

[9] M. F. Mammana, B. Micale, and M. Pennisi, Orthic quadrilaterals of a convex quadrilateral, *Forum Geom.*, 10 (2010) 79–91.

[10] L. J. Mordell and D. F. Barrow, (1937). Solution to Problem 3740, *Amer. Math. Monthly*, 44 (1937) 252–254.
[11] M. Şahin, *Matematik Olimpiyatlarına Hazırlık Geometri - I*, Palme Yayınları, Ankara, 2000.

Şahlar Meherrem: Yasar University, Faculty of Science and Letters, Department of Mathematics, Izmir, Turkey
E-mail address: sahlar.meherrem@yasar.edu.tr

Gizem Günel Açıksöz: Ozel Ege Lisesi, Izmir, Turkey
E-mail address: gizem.aciksoz@egelisesi.k12.tr

Serenay Şen : Ozel Ege Lisesi, Izmir, Turkey
E-mail address: serenay.sen3@gmail.com

Zeynep Sezer : Ozel Ege Lisesi, Izmir, Turkey
E-mail address: gunes_baskes@hotmail.com

Güneş Başkes: Ozel Ege Lisesi, Izmir, Turkey
E-mail address: zeynepsezer99@gmail.com