

## Sawayama and Thébault's theorem

Jean-Louis Ayme

**Abstract.** We present a purely synthetic proof of Thébault's theorem, known earlier to Y. Sawayama.

### 1. Introduction

In 1938 in a “Problems and Solutions” section of the *Monthly* [24], the famous French problemist Victor Thébault (1882-1960) proposed a problem about three circles with collinear centers (see Figure 1) to which he added a correct ratio and a relation which finally turned out to be wrong. The date of the first three metric




Figure 1

solutions [22] which appeared discreetly in 1973 in the Netherlands was more widely known in 1989 when the Canadian revue *Crux Mathematicorum* [27] published the simplified solution by Veldkamp who was one of the two first authors to prove the theorem in the Netherlands [26, 5, 6]. It was necessary to wait until the end of this same year when the Swiss R. Stark, a teacher of the Kantonsschule of Schaffhausen, published in the Helvetic revue *Elemente der Mathematik* [21] the first synthetic solution of a “more general problem” in which the one of Thébault's appeared as a particular case. This generalization, which gives a special importance to a rectangle known by J. Neuberg [15], citing [4], has been pointed out in 1983 by the editorial comment of the *Monthly* in an outline publication about the supposed

first metric solution of the English K. B. Taylor [23] which amounted to 24 pages. In 1986, a much shorter proof [25], due to Gerhard Turnwald, appeared. In 2001, R. Shail considered in his analytic approach, a “more complete” problem [19] in which the one of Stark appeared as a particular case. This last generalization was studied again by S. Gueron [11] in a metric and less complete way. In 2003, the *Monthly* published the angular solution by B. J. English, received in 1975 and “lost in the mists of time” [7].

Thanks to *JSTOR*, the present author has discovered in an ancient edition of the *Monthly* [18] that the problem of Shail was proposed in 1905 by an instructor Y. Sawayama of the central military School of Tokyo, and geometrically resolved by himself, mixing the synthetic and metric approach. On this basis, we elaborate a new, purely synthetic proof of Sawayama-Thébault theorem which includes several theorems that can all be synthetically proved. The initial step of our approach refers to the beginning of the Sawayama’s proof and the end refers to Stark’s proof. Furthermore, our point of view leads easily to the Sawayama-Shail result.

## 2. A lemma

**Lemma 1.** *Through the vertex  $A$  of a triangle  $ABC$ , a straight line  $AD$  is drawn, cutting the side  $BC$  at  $D$ . Let  $P$  be the center of the circle  $C_1$  which touches  $DC$ ,  $DA$  at  $E$ ,  $F$  and the circumcircle  $C_2$  of  $ABC$  at  $K$ . Then the chord of contact  $EF$  passes through the incenter  $I$  of triangle  $ABC$ .*




Figure 2

*Proof.* Let  $M$ ,  $N$  be the points of intersection of  $KE$ ,  $KF$  with  $C_2$ , and  $J$  the point of intersection of  $AM$  and  $EF$  (see Figure 3).  $KE$  is the internal bisector of  $\angle BKC$  [8, Théorème 119]. The point  $M$  being the midpoint of the arc  $BC$  which does not contain  $K$ ,  $AM$  is the  $A$ -internal bisector of  $ABC$  and passes through  $I$ .

The circles  $C_1$  and  $C_2$  being tangent at  $K$ ,  $EF$  and  $MN$  are parallel.




Figure 3

The circle  $C_2$ , the basic points  $A$  and  $K$ , the lines  $MAJ$  and  $NKF$ , the parallels  $MN$  and  $JF$ , lead to a converse of Reim's theorem ([8, Théorème 124]). Therefore, the points  $A$ ,  $K$ ,  $F$  and  $J$  are concyclic. This can also be seen directly from the fact that angles  $FJA$  and  $FKA$  are congruent.

Miquel's pivot theorem [14, 9] applied to the triangle  $AFJ$  by considering  $F$  on  $AF$ ,  $E$  on  $FJ$ , and  $J$  on  $AJ$ , shows that the circle  $C_4$  passing through  $E$ ,  $J$  and  $K$  is tangent to  $AJ$  at  $J$ . The circle  $C_5$  with center  $M$ , passing through  $B$ , also passes through  $I$  ([2, Livre II, p.46, théorème XXI] and [12, p.185]). This circle being orthogonal to circle  $C_1$  [13, 20] is also orthogonal to circle  $C_4$  ([10, 1]) as  $KEM$  is the radical axis of circles  $C_1$  and  $C_4$ .<sup>1</sup> Therefore,  $MB = MJ$ , and  $J = I$ . Conclusion: the chord of contact  $EF$  passes through the incenter  $I$ .  $\square$

*Remark.* When  $D$  is at  $B$ , this is the theorem of Nixon [16].

### 3. Sawayama-Thébault theorem

**Theorem 2.** *Through the vertex  $A$  of a triangle  $ABC$ , a straight line  $AD$  is drawn, cutting the side  $BC$  at  $D$ .  $I$  is the center of the incircle of triangle  $ABC$ . Let  $P$  be the center of the circle which touches  $DC$ ,  $DA$  at  $E$ ,  $F$ , and the circumcircle of  $ABC$ , and let  $Q$  be the center of a further circle which touches  $DB$ ,  $DA$  in  $G$ ,  $H$  and the circumcircle of  $ABC$ . Then  $P$ ,  $I$  and  $Q$  are collinear.*

<sup>1</sup>From  $\angle BKE = \angle MAC = \angle MBE$ , we see that the circumcircle of  $BKE$  is tangent to  $BM$  at  $B$ . So circle  $C_5$  is orthogonal to this circumcircle and consequently also to  $C_1$  as  $M$  lies on their radical axis.




Figure 4

*Proof.* According to the hypothesis,  $QG \perp BC$ ,  $BC \perp PE$ ; so  $QG \parallel PE$ . By Lemma 1,  $GH$  and  $EF$  pass through  $I$ . Triangles  $DHG$  and  $QGH$  being isosceles in  $D$  and  $Q$  respectively,  $DQ$  is

- (1) the perpendicular bisector of  $GH$ ,
- (2) the  $D$ -internal angle bisector of triangle  $DHG$ .

Mutatis mutandis,  $DP$  is

- (1) the perpendicular bisector of  $EF$ ,
- (2) the  $D$ -internal angle bisector of triangle  $DEF$ .

As the bisectors of two adjacent and supplementary angles are perpendicular, we have  $DQ \perp DP$ . Therefore,  $GH \parallel DP$  and  $DQ \parallel EF$ . Conclusion: using the converse of Pappus's theorem ([17, Proposition 139] and [3, p.67]), applied to the hexagon  $PEIGQDP$ , the points  $P$ ,  $I$  and  $Q$  are collinear.  $\square$

## References

- [1] N. Altshiller-Court, *College Geometry*, Barnes & Noble, 205.
- [2] E. Catalan, *Théorèmes et problèmes de Géométrie élémentaires*, 1879.
- [3] H. S. M. Coxeter and S. L. Greitzer, *Geometry Revisited*, Math. Assoc. America, 1967.
- [4] *Archiv der Mathematik und Physik* (1842) 328.
- [5] B. C. Dijkstra-Kluyver, Twee oude vraagstukken in één klap opgelost, *Nieuw Tijdschrift voor Wiskunde*, 61 (1973-74) 134–135.
- [6] B. C. Dijkstra-Kluyver and H. Streefkerk, Nogmaals het vraagstuk van Thébault, *Nieuw Tijdschrift voor Wiskunde*, 61 (1973-74) 172–173.
- [7] B. J. English, Solution of Problem 3887, *Amer. Math. Monthly*, 110 (2003) 156–158.
- [8] F. G.-M., *Exercices de Géométrie*, sixième édition, 1920, J. Gabay reprint.
- [9] H. G. Forder, *Geometry*, Hutchinson, 1960.
- [10] L. Gaultier (de Tours), Les contacts des cercles, *Journal de l'Ecole Polytechnique*, Cahier 16 (1813) 124–214.
- [11] S. Gueron, Two Applications of the Generalized Ptolemy Theorem, *Amer. Math. Monthly*, 109 (2002) 362–370.

- [12] R. A. Johnson, *Advanced Euclidean Geometry*, Dover, 1965.
- [13] *Leybourn's Mathematical Repository* (Nouvelle série) 6 tome I, 209.
- [14] A. Miquel, Théorèmes de Géométrie, *Journal de mathématiques pures et appliquées de Liouville*, 3 (1838) 485-487.
- [15] J. Neuberg, *Nouvelle correspondance mathématique*, 1 (1874) 96.
- [16] R. C. J. Nixon, Question 10693, *Reprints of Educational Times*, London (1863-1918) 55 (1891) 107.
- [17] Pappus, *La collection mathématique*, 2 volumes, French translation by Paul Ver Eecker, Paris, Desclée de Brouwer, 1933.
- [18] Y. Sawayama, A new geometrical proposition, *Amer. Math. Monthly*, 12 (1905) 222-224.
- [19] R. Shail., A proof of Thébault's Theorem, *Amer. Math. Monthly*, 108 (2001) 319-325.
- [20] S. Shirali, On the generalized Ptolemy theorem, *Crux Math.*, 22 (1996) 48-53.
- [21] R. Stark, Eine weitere Lösung der Thébault'schen Aufgabe, *Elem. Math.*, 44 (1989) 130-133.
- [22] H. Streefkerk, Waarom eenvoudig als het ook ingewikkeld kan?, *Nieuw Tijdschrift voor Wiskunde*, 60 (1972-73) 240-253.
- [23] K. B. Taylor, Solution of Problem 3887, *Amer. Math. Monthly*, 90 (1983) 482-487.
- [24] V. Thébault, Problem 3887, Three circles with collinear centers, *Amer. Math. Monthly*, 45 (1938) 482-483.
- [25] G. Turnwald, Über eine Vermutung von Thébault, *Elem. Math.*, 41 (1986) 11-13.
- [26] G. R. Veldkamp, Een vraagstuk van Thébault uit 1938, *Nieuw Tijdschrift voor Wiskunde*, 61 (1973-74) 86-89.
- [27] G. R. Veldkamp, Solution to Problem 1260, *Crux Math.*, 15 (1989) 51-53.

Jean-Louis Ayme: 37 rue Ste-Marie, 97400 St.-Denis, La Réunion, France  
E-mail address: jeanlouisayme@yahoo.fr