

Some More Archimedean Circles in the Arbelos

Frank Power

Abstract. We construct 4 circles in the arbelos which are congruent to the Archimedean twin circles.

Thomas Schoch [2] tells the remarkable story of his discovery in the 1970's of the many Archimedean circles in the arbelos (shoemaker's knife) that were eventually recorded in the paper [1]. In this note, we record four more Archimedean circles which were discovered in the summer of 1998, when the present author took a geometry course ([3]) with one of the authors of [1].

Consider an arbelos with inner semicircles C_1 and C_2 of radii a and b , and outer semicircle C of radius $a + b$. It is known the Archimedean circles have radius $t = \frac{ab}{a+b}$. Let Q_1 and Q_2 be the "highest" points of C_1 and C_2 respectively.

Theorem. A circle tangent to C internally and to OQ_1 at Q_1 (or OQ_2 at Q_2) has radius $t = \frac{ab}{a+b}$.




Figure 1

Proof. There are two such circles tangent at Q_1 , namely, (C_1) and (C'_1) in Figure 1. Consider one such circle (C_1) with radius r . Note that

$$OQ_1^2 = O_1Q_1^2 + OO_1^2 = a^2 + b^2.$$

It follows that

$$(a + b - r)^2 = (a^2 + b^2) + r^2,$$

from which $r = \frac{ab}{a+b} = t$. The same calculation shows that (C'_1) also has radius t , and similarly for the two circles at Q_2 . \square

References

- [1] C. W. Dodge, T. Schoch, P. Y. Woo and P. Yiu, Those ubiquitous Archimedean circles, *Math. Mag.*, 72 (1999) 202–213.
- [2] T. Schoch, Arbelos, <http://www.retas.de/thomas/arbelos/arbelos.html>.
- [3] P. Yiu, *Euclidean Geometry*, available at <http://www.math.fau.edu/Yiu/Geometry.html>.

Frank Power: Atlantic Community High School, 2455 W. Atlantic Avenue, Delray Beach, Florida, 33445, USA

E-mail address: `powerrf@palmbeach.k12.fl.us`